

Recombinant Human IL-23 (Interleukin 23)

Catalog # (Size): HZ-1254 (10µg) HZ-1261 (100µg) HZ-1262 (1000µg)

Product Description

- Endotoxin-free
- Animal-derived product free
- Available in Bulk
- Lyophilized and Carrier Free (CF)
- Expressed as a disulfide-linked heterodimer

Xeno-free IL-23 ^{HuXp} is expressed in human 293 cells as a heterodimeric glycoprotein composed of two disulfide-linked subunits (p40 cysteine linked to p19). IL-23 has been shown to enhance proliferation of memory T cells. It also stimulates the production of IFN-gamma in NK cells, induces IL-17 production, and drives Th17 mediated responses. Furthermore, it is known that IL-23 takes a vital part in the inflammation process and that it is associated with auto immune diseases.

Typical Specifications

Species	Human
Expression	HEK293 Cell Expressed
Activity	Typically ≤ 30 ng/mL EC ₅₀
Purity	>95%
Endotoxin	<1 EU/µg
Molecular Mass	55 kDa, heterodimer, glycosylated
Formulation	1x PBS

Purity Confirmation



The protein was resolved by SDS-polyacrylamide gel electrophoresis and the gel was stained with Coomassie blue.

Activity Assay

The activity was determined by the dose-dependent secretion of IL-17 in response to IL-23 using murine splenocytes.



All HumaXpress® HumanKine™ are animal-component-free and Xeno-free™



Reconstitution Buffer

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 1x PBS containing 0.1% endotoxin-free recombinant human serum albumin (HSA).

Limited Use and Restrictions Unless otherwise stated in our catalog or other company documentation accompanying the products sold by HumanZyme Inc. are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, including resale or use in manufacture, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.