

**Recombinant Human
TGF beta 2 (Transforming Growth Factor beta 2)**

Catalog # (Size): HZ-1092 (5µg) HZ-1199 (100µg) HZ-1088 (1000µg)

Product Description

- Endotoxin-free
- Animal-derived product free
- Available in Bulk
- Lyophilized and Carrier Free (CF)
- High Activity

Xeno-free TGF beta 2^{HuXp} is expressed from human 293 cells as a mature disulfide-linked homodimer with apparent molecular mass of 25 kDa. This cytokine is produced in a human cell expression system with serum-free, chemically defined media. The cytokine is greater than 95% pure.

Typical Specifications

Species	Human
Expression	HEK293 Cell Expressed
Activity	Typically ≤ 0.5 ng/mL EC ₅₀
Purity	>95%
Endotoxin	<1 EU/µg
Molecular Mass	25 kDa, homodimer, non-glycosylated
Formulation	50 mM NaOAc pH 4.5

Purity Confirmation



The protein was resolved by SDS-polyacrylamide gel electrophoresis and the gel was stained with Coomassie blue.

Activity Assay

The activity was determined by the dose dependent inhibition of IL-4 induced proliferation of mouse HT-2 cells (BALB/c spleen activated by sheep erythrocytes in the presence of IL-2).



All HumaXpress® HumanKine™ are animal-component-free and Xeno-free™



Reconstitution Buffer

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 20% ethanol + 50 mM NaOAc + 75 mM HOAc.

Limited Use and Restrictions Unless otherwise stated in our catalog or other company documentation accompanying the products sold by HumanZyme Inc. are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, including resale or use in manufacture, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.